


# BPH SYMPTOM RELIEF THAT WORKS NIGHTS SO HE CAN WORK DAYS



## WELCOME TO THE RAPAFLO<sup>®</sup> PROFESSIONAL SITE

RAPAFLO<sup>®</sup> is a uniquely selective alpha-blocker that provides rapid and sustained relief of benign prostatic hyperplasia (BPH) symptoms with a low incidence of vasodilatory and orthostatic effects.<sup>1-3</sup> RAPAFLO<sup>®</sup> has been proven safe and effective for the treatment of frequency, nocturia, and other bothersome BPH symptoms in multiple patient types.<sup>2,3</sup>

### AN ONLINE RESOURCE FOR HEALTH CARE PROFESSIONALS TO LEARN ABOUT:

The safety, efficacy, and mechanism of action of [RAPAFLO<sup>®</sup>](#)

RAPAFLO<sup>®</sup> [dosing and administration](#)

RAPAFLO<sup>®</sup> [access and reimbursement](#)

[How to start a conversation](#) about BPH with your older male patients

How to [assess the severity](#) of your patients' BPH symptoms and make a diagnosis

[BPH and its impact](#)

Appropriate [treatment options](#) for BPH

[Professional resources](#) for you and your patients

### + REFERENCES

RAPAFLO<sup>®</sup> is indicated for the treatment of the signs and symptoms of benign prostatic hyperplasia (BPH). RAPAFLO<sup>®</sup> is not indicated for the treatment of hypertension.

### IMPORTANT SAFETY INFORMATION

RAPAFLO<sup>®</sup> is contraindicated in patients with severe renal impairment (CCr <30 mL/min), severe hepatic impairment (Child-Pugh score ≥10), with use of strong CYP3A4 inhibitors, and in patients with a history of hypersensitivity to silodosin or any of the ingredients of RAPAFLO<sup>®</sup>.

Postural hypotension with or without symptoms (eg, dizziness) may develop when beginning treatment with RAPAFLO<sup>®</sup>. As with all alpha-blockers, there is a potential for syncope. Patients should be warned of the possible occurrences of such events and should avoid situations where injury could result. RAPAFLO<sup>®</sup> should be used with caution in patients with moderate renal impairment. Patients should be assessed to rule out the presence of prostate cancer prior to starting treatment with RAPAFLO<sup>®</sup>. Patients planning cataract surgery should inform their ophthalmologist that they are taking RAPAFLO<sup>®</sup>.

The most common side effects are retrograde ejaculation, dizziness, diarrhea, orthostatic hypotension, headache, nasopharyngitis, and nasal congestion.



Models are for illustrative purposes only.

Copyright © 2008-2013 Actavis, Inc. All rights reserved.

[LEGAL/COPYRIGHT DISCLAIMER](#) | [PRIVACY POLICY](#) | 09505 | 6/13